

4-Hydroxymethyl, 4-Methyl and 4-Hydroxy Benzenediazonium and Their Salts

- 4-Hydroxymethyl benzenediazonium (4-HMBD) is naturally present in the commonly cultivated edible mushroom *Agaricus bisporus*.
- 4-Methyl benzenediazonium (4-MBD) may form from 4-HMBD.
- 4-Hydroxy benzenediazonium (4-HBD) is naturally present in *Agaricus xanthodermus*, a wild inedible mushroom that may be mistaken for *Agaricus bisporus*.

Consumption of *Agaricus bisporus* mushrooms may result in exposure to 4-HMBD and its salts, and possibly, to 4-MBD and its salts. If *Agaricus xanthodermus* mushrooms are mistakenly ingested, exposure to 4-HBD and its salts may occur.

These benzenediazonium compounds and their salts each passed the animal data screen, underwent a preliminary toxicological evaluation, and are being brought to the Carcinogen Identification Committee for consultation. This is a compilation of the relevant studies identified during the preliminary toxicological evaluation.

Epidemiological data

No cancer epidemiology studies on these benzenediazonium compounds were identified.

Animal carcinogenicity data

4-HMBD

- Subcutaneous injection studies of 4-HMBD tetrafluoroborate
 - Male and female Swiss mice (injected once per week for 26 weeks, observed for life): Toth *et al.* (1981)
 - *Increases in tumors of the subcutis and skin in females and males (by pairwise comparisons)*
- Gavage studies of 4-HMBD tetrafluoroborate
 - Male and female Swiss mice (single gavage, observed for life): Toth *et al.* (1982)
 - *Increases in tumors of the glandular stomach in females and males (by pairwise comparisons)*

- Subcutaneous injection studies of 4-HMBD sulfate
 - Male and female Swiss mice (injected once per week for 26 weeks, observed for life): Toth (1987)
 - *Increases in tumors of the subcutis and skin in females and males (by pairwise comparisons)*

4-MBD

- Subcutaneous injection studies of 4-MBD sulfate
 - Male and female Swiss mice (injected once per week for 16 to 19 weeks, observed for life): Toth *et al.* (1989b)
 - *Increases in tumors of the subcutis and skin in females (by pairwise comparisons) and of the subcutis in males (by pairwise comparison)*

4-HBD

- Subcutaneous injection studies of 4-HBD sulfate
 - Male and female Swiss mice (injected once per week for 36 weeks, observed for life): Toth *et al.* (1989a)
 - *Increases in tumors of the subcutis in females and males (by pairwise comparisons)*
 - Male and female Swiss mice (injected once per week for 2 weeks, observed for life): Toth *et al.* (1989a)
 - *No treatment-related tumor findings*
- Five-week skin painting study of 4-HBD
 - Male Wistar rats \pm magnesium-deficient diet: Quintero *et al.* (2006)
 - *Increase in incipient fibrosarcoma in treated magnesium-deficient vs. untreated magnesium-deficient animals (and vs. treated magnesium-sufficient animals)*

Other relevant data

- Genotoxicity
 - 4-HMBD
 - *In vitro* DNA damage/base modification, single strand breaks (*positive*): Lawson *et al.* (1995); Hiramoto *et al.* (1995a, 1995b)
 - *In vivo* mouse liver DNA adduct formation (*positive*): Hiramoto *et al.* (1998)
 - Mutations in *Salmonella* (*positive*): Lawson *et al.* (1995)
 - Mutations in V79 cells (*weakly positive*): Lawson *et al.* (1995)
 - DNA cross-links in V79 cells (*positive*): Lawson *et al.* (1995)

- *In vivo* mouse peripheral reticulocyte micronuclei (*positive*): Hiramoto *et al.* (1995b)
 - 4-MBD
 - *In vitro* DNA damage/base modification, single strand breaks (*positive*): Lawson *et al.* (1995)
 - Mutations in *Salmonella* (*positive*): Lawson *et al.* (1995)
 - Mutations in V79 cells (*weakly positive*): Lawson *et al.* (1995)
 - DNA cross-links in V79 cells (*negative*): Lawson *et al.* (1995)
 - DNA adduct formation in the mouse C50 keratinocyte cell line and mouse primary keratinocytes (*positive*): Gannett *et al.* (1996)
- Mechanistic considerations
 - 4-MBD activation of transcription factor AP-1 in a mouse cell line and in mice *in vivo* (*positive*): Gannett *et al.* (2000)
 - Review of mechanistic information on arenediazonium ion carcinogenicity: Powell and Gannett (2002)
- Structure activity considerations
 - 4-HMBD, 4-HBD, and 4-MBD are structurally similar to each other, and all three induce skin and subcutaneous tumors in male and female mice
 - A fourth structurally related compound, benzenediazonium sulphate, also induces skin and subcutaneous tumors in male and female mice: Toth *et al.* (1999)

References¹

Gannett PM, Lawson T, Miller M, Thakkar DD, Lord JW, Yau WM, Toth B (1996). 8-Arylguanine adducts from arenediazonium ions and DNA. *Chem Biol Interact* **101**:149-64.

Gannett PM, Ye J, Ding M, Powell J, Zhang Y, Darian E, Daft J, Shi X (2000). Activation of AP-1 through the MAP kinase pathway: a potential mechanism of the carcinogenic effect of arenediazonium ions. *Chem Res Toxicol* **13**:1020-7.

Hiramoto K, Kaku M, Sueyoshi A, Fujise M, Kikugawa K (1995a). DNA base and deoxyribose modification by the carbon-centered radical generated from

¹ Excerpts or the complete publication have been provided to members of the Carcinogen Identification Committee, in the order in which they are discussed in this document.

4-(hydroxymethyl) benzenediazonium salt, a carcinogen in mushroom. *Chem Res & Toxi* **8**:356-62.

Hiramoto K, Kaku M, Kato T, Kikugawa K (1995b). DNA strand breaking by the carbon-centered radical generated from 4-(hydroxymethyl) benzenediazonium salt, a carcinogen in mushroom *Agaricus bisporus*. *Chem Bio Interact* **94**:21-36.

Hiramoto K, Ando M, Kato T, Kikugawa K (1998). ³²P-postlabeling detection of DNA base adducts in mice induced by 4-(hydroxymethyl)benzenediazonium salt, a carcinogen in mushroom *Agaricus bisporus*. *Japn J Toxi & Env Health* **44**:300-304.

Lawson T, Gannett PM, Yau W, Dalal NS, Toth B (1995). Different patterns of mutagenicity of arenediazonium ions in V79 cells and *Salmonella typhimurium* TA102: Evidence for different mechanisms of action. *J Agri Food Chem* **43**:2627-35.

Powell JH, Gannett PM (2002). Mechanisms of carcinogenicity of aryl hydrazines, aryl hydrazides, and arenediazonium ions. *J Environ Pathol Toxi Oncol* **21**:1-31.

Quintero B, Planells E, del Cabeza Cabeza M, Esquivias J, del Pilar Gutiérrez M, Sánchez C, Aranda P, Zarzuelo A, Llopis J (2006). Tumor promoting activity of *p*-hydroxybenzenediazonium is accelerated in Mg-deficient rats. *Chem Biol Interact* **159**:186-195.

Toth B (1987). Cancer induction by the sulfate form of 4-(hydroxymethyl) benzenediazonium ion of *Agaricus bisporus*. *In Vivo* **1**(1):39-42.

Toth B, Nagel D, Ross A (1982). Gastric tumorigenesis by a single dose of 4-(hydroxymethyl)benzenediazonium ion of *Agaricus bisporus*. *Br J Cancer* **46**(3):417-22.

Toth B, Patil K, Erickson J, Gannett P (1999). Cancer induction studies using different administrations of benzenediazonium sulphate in mice. *In Vivo* **13**:125-128.

Toth B, Patil K, Jae H-S (1981). Carcinogenesis of 4-(hydroxymethyl) benzenediazonium ion (tetrafluoroborate) of *Agaricus bisporus*. *Cancer Res* **41**(6):2444-9.

Toth B, Patil K, Taylor J, Stessman C, Gannett P (1989a). Cancer induction in mice by 4-hydroxybenzenediazonium sulfate of the *Agaricus xanthodermus* mushroom. *In Vivo* **3**(5):301-6.

Toth B, Taylor J, Mattson B, Gannett P (1989b). Tumor induction by 4-(methyl)benzenediazonium sulfate in mice. *In Vivo* **3**(1):17-21.